


Stoma Site Selection

Stoma site selection and marking should be done for all patients scheduled for ostomy surgery by an experienced, educated, competent clinician. This should be considered even if there is only a possibility for stoma creation. For a pouch to fit comfortably and securely, it is important to have an adequate, intact skin surface. Proper placement helps prevent skin and stoma complications, pouching problems, pain, and clothing concerns. The optimal site enhances the likelihood of independence in stoma care and resumption of normal activities. The preoperative visit also provides an opportunity for education for patient and family.

Anatomy



IDEAL STOMA CHARACTERISTICS:	SITES TO AVOID:	OTHER CHARACTERISTICS:
• Red • Round • Raised (about 1" protrusion) • Lumen at center of stoma • Smooth skin surface	• Scars/Wrinkles/Incision Lines • Skin folds/Creases • Bony prominence • Under pendulous breasts • Suture lines • Umbilicus • Belt/Waistline • Hernia • Mobile abdominal tissue • Radiation sites	• Type of ostomy • Occupation • Impairments (e.g., visual, physical) • Sports/Activity level • Prosthetic equipment • Location preference (surgeon, patient) • Multiple stoma sites (consider differing levels) • Age • Diagnosis • Posture • Contractures

Stoma Site Selection

To select the correct site for the stoma, the following assessments should be made preoperatively:

Type of stoma anticipated	The abdomen can be divided into four quadrants. Correlating this topographical information to the underlying anatomical structures will help locate the correct quadrant for the stoma. For example, an ileostomy (ileum) would usually be located in the right lower quadrant.
Rectus muscle sheath	Placement in the rectus muscle can help prevent some stomal complications. This muscle runs vertically through the abdomen (refer to diagram) and may be located by inspection and/or palpation.
Adequate surface area	The pouching system is secured by adhesive. There needs to be an adequate adhesive contact surface between the pouch and the skin for secure attachment. Generally an area of two to three inches of flat surface is preferred but not always possible. On a child, a smaller area is needed and is dependent on body size. A stoma siting disc can be used to evaluate adequate skin surface area preoperatively.
Easily seen	It is difficult for a person to be independent in their care if they cannot see their stoma. Select a site visible to the patient and if possible, below the belt line to conceal the pouch. For many people, the best location is in the lower quadrant on the apex of the infraumbilical bulge. If the patient is extremely obese, place mark in upper quadrant.
Smooth skin surface	Locating the stoma in an area where the skin is flat is important. The prospective stoma site should be located away from skin folds, bony prominence, scars, the umbilicus, incision lines, and the belt line. Any of these can interfere with a secure pouch seal. The abdomen should be observed in various positions. Refer to diagram to the right and sites to avoid on front.
Miscellaneous criteria	In addition there may be other factors that could impact where the stoma would ideally be located; for example, if a person uses a wheelchair, walker, or brace, or needs more than one stoma. Fecal and urinary stomas should be marked on different horizontal planes/lines. Refer to front for other considerations.

Positions

Evaluate potential site in lying, sitting, bending, and standing positions.



Lying



Sitting



Bending



Standing

Marking the Site

After the site is selected, it needs to be marked. The technique used for marking the site varies and may be based on hospital protocols. The site location should be documented and communicated to the surgeon so that the intent of the mark is understood in the operating room. An indelible marker or skin dye may be used to identify the site but must be visible after the surgical scrub. In some cases, two choices may be made with the first labeled as #1. Preoperative site markings are a guide. Final selection is done by the surgeon during the operative procedure.

For product questions, sampling needs, or detailed clinical questions concerning our products in the US, call **1.888.808.7456**. In Canada call **1.800.263.7400**.

Routine follow-up with your healthcare professional is recommended.

Prior to using any ostomy products/accessories be sure to read all product inserts and labels.

Wound, Ostomy and Continence Nurses Society. (2014). *WOCN Society and ASCRS Position Statement on Preoperative Stoma Site Marking for Patients Undergoing Colostomy or Ileostomy Surgery*. Mt. Laurel: NJ. Stoma Siting Procedure. www.wocn.org J.E. Carmel, J.C. Colwell, M.T. Goldberg (Eds.), *WOCN Society Core Curriculum Ostomy Management* (Chapter 8). Philadelphia: Wolters Kluwer. 2016.

The Hollister logo and "Healthy skin. Positive outcomes." are trademarks of Hollister Incorporated.
© 2017 Hollister Incorporated. 923080-1217 US-00228

Hollister Incorporated
2000 Hollister Drive
Libertyville, Illinois 60048
1.800.323.4060

www.hollister.com

Hollister Limited
95 Mary Street
Aurora, Ontario L4G 1G3
1.800.263.7400